

BILL, RECORD LECTURE!!!!

BILL RECORD LECTURE!!!

The Distinct Volumes Problem

David Conlon- Cambridge (Prof)

Jacob Fox-MIT (Prof)

William Gasarch-U of MD (Prof)

David Harris- U of MD (Grad Student)

Douglas Ulrich- U of MD (Ugrad Student)

Sam Zbarsky- Mont. Blair. (High School Student- now CMU)

Darling Wants an Actual Coloring

1. **Infinite Ramsey Thm:** For any 2-coloring of the EDGES of K_ω there exists an infinite *monochromatic* K_ω .

Darling Wants an Actual Coloring

1. **Infinite Ramsey Thm:** For any 2-coloring of the EDGES of K_ω there exists an infinite *monochromatic* K_ω .
2. **Infinite Can Ramsey Thm:** For any ω -coloring of the EDGES of K_ω there exists an infinite H such that either (1) H homog, (2) H min-homog, (3) H max-homog, (4) H rainbow.

Darling Wants an Actual Coloring

1. **Infinite Ramsey Thm:** For any 2-coloring of the EDGES of K_ω there exists an infinite *monochromatic* K_ω .
2. **Infinite Can Ramsey Thm:** For any ω -coloring of the EDGES of K_ω there exists an infinite H such that either (1) H homog, (2) H min-homog, (3) H max-homog, (4) H rainbow.
3. **Darling:** Give me an **example** of an actual coloring.

Darling Wants an Actual Coloring

1. **Infinite Ramsey Thm:** For any 2-coloring of the EDGES of K_ω there exists an infinite *monochromatic* K_ω .
2. **Infinite Can Ramsey Thm:** For any ω -coloring of the EDGES of K_ω there exists an infinite H such that either (1) H homog, (2) H min-homog, (3) H max-homog, (4) H rainbow.
3. **Darling:** Give me an **example** of an actual coloring.

Bill thinks of one— next page.

Points and Distances

Let p_1, p_2, \dots be an infinite set of points in \mathbb{R} .

Points and Distances

Let p_1, p_2, \dots be an infinite set of points in \mathbb{R} .

Let $COL : \binom{\mathbb{N}}{2} \rightarrow \mathbb{R}$ be

$$COL(i, j) = |p_i - p_j| \text{ (Distance Between } p_i \text{ and } p_j).$$

Points and Distances

Let p_1, p_2, \dots be an infinite set of points in \mathbb{R} .

Let $COL : \binom{\mathbb{N}}{2} \rightarrow \mathbb{R}$ be

$$COL(i, j) = |p_i - p_j| \text{ (Distance Between } p_i \text{ and } p_j).$$

Result: For any infinite set of points in the plane there is an infinite subset where all distances are distinct. (Already known by Erdős via diff proof.)

Points and Distances

Let p_1, p_2, \dots be an infinite set of points in \mathbb{R}^2 .

Let $COL : \binom{\mathbb{N}}{2} \rightarrow \mathbb{R}$ be

$$COL(i, j) = |p_i - p_j| \text{ (Distance Between } p_i \text{ and } p_j).$$

Result: For any infinite set of points in the plane there is an infinite subset where all distances are distinct. (Already known by Erdős via diff proof.)

Next Step: Finite version: Can use Finite Can Ramsey to prove the following: For every set of n points in the plane there is a subset of size $\Omega(\log n)$ where all distances are distinct. (Much better is known.)

INITIAL MOTIVATION ABANDONED

INITIAL MOTIVATION ABANDONED

1. Dumped Ramsey! Added co-authors! Got **new** results!

INITIAL MOTIVATION ABANDONED

1. Dumped Ramsey! Added co-authors! Got **new** results!
2. What about **Area**? If there are n points in \mathbb{R}^2 want large subset so that all areas are distinct.

INITIAL MOTIVATION ABANDONED

1. Dumped Ramsey! Added co-authors! Got **new** results!
2. What about **Area**? If there are n points in \mathbb{R}^2 want large subset so that all areas are distinct.
3. More general question: n points in \mathbb{R}^d and looking for all a -volumes to be different. (This question seems to be new.)

EXAMPLES with DISTANCES

The following is an **EXAMPLE** of the kind of theorems we will be talking about.

*If there are n points in \mathbb{R}^2 then there is a subset of size $\Omega(n^{1/3})$ with all distances between points **DIFF**.*

EXAMPLES with AREAS

*If there are n points in \mathbb{R}^2 then there is a subset of size $\Omega(n^{1/5})$ with all triangle areas **DIFF**.*

EXAMPLES with AREAS

*If there are n points in \mathbb{R}^2 then there is a subset of size $\Omega(n^{1/5})$ with all triangle areas **DIFF**.*

FALSE: Take n points on a LINE. All triangle areas are 0.

EXAMPLES with AREAS

*If there are n points in \mathbb{R}^2 then there is a subset of size $\Omega(n^{1/5})$ with all triangle areas **DIFF**.*

FALSE: Take n points on a LINE. All triangle areas are 0.

We state theorems in **no three collinear** form to get around this issue.

Maximal Rainbow Sets

Def: A **(2)-Rainbow Set** is a set of points in \mathbb{R}^d where all of the distances are distinct. Also called a **dist-rainbow**.

Maximal Rainbow Sets

Def: A **(2)-Rainbow Set** is a set of points in \mathbb{R}^d where all of the distances are distinct. Also called a **dist-rainbow**.

Def: A **3-Rainbow Set** is a set of points in \mathbb{R}^d where all nonzero areas of triangles are distinct. Also called an **area-rainbow**.

Maximal Rainbow Sets

Def: A **(2)-Rainbow Set** is a set of points in \mathbb{R}^d where all of the distances are distinct. Also called a **dist-rainbow**.

Def: A **3-Rainbow Set** is a set of points in \mathbb{R}^d where all nonzero areas of triangles are distinct. Also called an **area-rainbow**.

Def: An **a -Rainbow Set** is a set of points in \mathbb{R}^d where all nonzero a -volumes are distinct. An a -volume is the volume enclosed by a points. Also called a **vol-rainbow**.

Maximal Rainbow Sets

Def: A **(2)-Rainbow Set** is a set of points in \mathbb{R}^d where all of the distances are distinct. Also called a **dist-rainbow**.

Def: A **3-Rainbow Set** is a set of points in \mathbb{R}^d where all nonzero areas of triangles are distinct. Also called an **area-rainbow**.

Def: An **a -Rainbow Set** is a set of points in \mathbb{R}^d where all nonzero a -volumes are distinct. An a -volume is the volume enclosed by a points. Also called a **vol-rainbow**.

Def: Let $X \subseteq \mathbb{R}^d$. A **Maximal Rainbow Set** is a rainbow set $Y \subseteq X$ such that if any more points of X are added then it STOPS being a rainbow set.

Maximal Rainbow Sets

Def: A **(2)-Rainbow Set** is a set of points in \mathbb{R}^d where all of the distances are distinct. Also called a **dist-rainbow**.

Def: A **3-Rainbow Set** is a set of points in \mathbb{R}^d where all nonzero areas of triangles are distinct. Also called an **area-rainbow**.

Def: An **a -Rainbow Set** is a set of points in \mathbb{R}^d where all nonzero a -volumes are distinct. An a -volume is the volume enclosed by a points. Also called a **vol-rainbow**.

Def: Let $X \subseteq \mathbb{R}^d$. A **Maximal Rainbow Set** is a rainbow set $Y \subseteq X$ such that if any more points of X are added then it STOPS being a rainbow set.

Def: Let $X \subseteq \mathbb{R}^d$. An **a -Maximal Rainbow Set** is a a -rainbow set $Y \subseteq X$ such that if any more points of X are added then it STOPS being an a -rainbow set.

Easy Lemma

Lemma If there is a MAP from X to Y that is $\leq c$ -to-1 then
 $|Y| \geq |X|/c$.

We will call this LEMMA.

The $d = 1$ Case

Thm: For all $X \subseteq \mathbb{R}^1$ of size n there exists a dist-rainbow subset of size $\Omega(n^{1/3})$.

The $d = 1$ Case

Thm: For all $X \subseteq \mathbb{R}^1$ of size n there exists a dist-rainbow subset of size $\Omega(n^{1/3})$.

Proof: Let M be a **MAXIMAL DIST-RAINBOW SET**.
Let $x \in X - M$. WHY IS x NOT IN M !? Either

The $d = 1$ Case

Thm: For all $X \subseteq \mathbb{R}^1$ of size n there exists a dist-rainbow subset of size $\Omega(n^{1/3})$.

Proof: Let M be a **MAXIMAL DIST-RAINBOW SET**.

Let $x \in X - M$. WHY IS x NOT IN M !? Either

- ▶ $(\exists x_1, x_2 \in M)[|x - x_1| = |x - x_2|]$, or
- ▶ $(\exists x_1, x_2, x_3 \in M)[|x - x_1| = |x_2 - x_3|]$.

The $d = 1$ Case

Thm: For all $X \subseteq \mathbb{R}^1$ of size n there exists a dist-rainbow subset of size $\Omega(n^{1/3})$.

Proof: Let M be a **MAXIMAL DIST-RAINBOW SET**.

Let $x \in X - M$. WHY IS x NOT IN M !? Either

- ▶ $(\exists x_1, x_2 \in M)[|x - x_1| = |x - x_2|]$, or
- ▶ $(\exists x_1, x_2, x_3 \in M)[|x - x_1| = |x_2 - x_3|]$.

f maps an element of $X - M$ to **reason** $x \notin M$.

The $d = 1$ Case

Thm: For all $X \subseteq \mathbb{R}^1$ of size n there exists a dist-rainbow subset of size $\Omega(n^{1/3})$.

Proof: Let M be a **MAXIMAL DIST-RAINBOW SET**.

Let $x \in X - M$. WHY IS x NOT IN M !? Either

- ▶ $(\exists x_1, x_2 \in M)[|x - x_1| = |x - x_2|]$, or
- ▶ $(\exists x_1, x_2, x_3 \in M)[|x - x_1| = |x_2 - x_3|]$.

f maps an element of $X - M$ to **reason** $x \notin M$.

$$f : X - M \rightarrow \binom{M}{2} \cup M \times \binom{M}{2}$$

The $d = 1$ Case

Thm: For all $X \subseteq \mathbb{R}^1$ of size n there exists a dist-rainbow subset of size $\Omega(n^{1/3})$.

Proof: Let M be a **MAXIMAL DIST-RAINBOW SET**.

Let $x \in X - M$. WHY IS x NOT IN M !? Either

- ▶ $(\exists x_1, x_2 \in M)[|x - x_1| = |x - x_2|]$, or
- ▶ $(\exists x_1, x_2, x_3 \in M)[|x - x_1| = |x_2 - x_3|]$.

f maps an element of $X - M$ to **reason** $x \notin M$.

$$f : X - M \rightarrow \binom{M}{2} \cup M \times \binom{M}{2}$$

What is $f^{-1}(\{x_1, x_2\})$?

The $d = 1$ Case

Thm: For all $X \subseteq \mathbb{R}^1$ of size n there exists a dist-rainbow subset of size $\Omega(n^{1/3})$.

Proof: Let M be a **MAXIMAL DIST-RAINBOW SET**.

Let $x \in X - M$. WHY IS x NOT IN M !? Either

- ▶ $(\exists x_1, x_2 \in M)[|x - x_1| = |x - x_2|]$, or
- ▶ $(\exists x_1, x_2, x_3 \in M)[|x - x_1| = |x_2 - x_3|]$.

f maps an element of $X - M$ to **reason** $x \notin M$.

$$f : X - M \rightarrow \binom{M}{2} \cup M \times \binom{M}{2}$$

What is $f^{-1}(\{x_1, x_2\})$? It's ≤ 1 POINT.

The $d = 1$ Case

Thm: For all $X \subseteq \mathbb{R}^1$ of size n there exists a dist-rainbow subset of size $\Omega(n^{1/3})$.

Proof: Let M be a **MAXIMAL DIST-RAINBOW SET**.

Let $x \in X - M$. WHY IS x NOT IN M !? Either

- ▶ $(\exists x_1, x_2 \in M)[|x - x_1| = |x - x_2|]$, or
- ▶ $(\exists x_1, x_2, x_3 \in M)[|x - x_1| = |x_2 - x_3|]$.

f maps an element of $X - M$ to **reason** $x \notin M$.

$$f : X - M \rightarrow \binom{M}{2} \cup M \times \binom{M}{2}$$

What is $f^{-1}(\{x_1, x_2\})$? It's ≤ 1 POINT.

What is $f^{-1}(x_1, \{x_2, x_3\})$?

The $d = 1$ Case

Thm: For all $X \subseteq \mathbb{R}^1$ of size n there exists a dist-rainbow subset of size $\Omega(n^{1/3})$.

Proof: Let M be a **MAXIMAL DIST-RAINBOW SET**.

Let $x \in X - M$. WHY IS x NOT IN M !? Either

- ▶ $(\exists x_1, x_2 \in M)[|x - x_1| = |x - x_2|]$, or
- ▶ $(\exists x_1, x_2, x_3 \in M)[|x - x_1| = |x_2 - x_3|]$.

f maps an element of $X - M$ to **reason** $x \notin M$.

$$f : X - M \rightarrow \binom{M}{2} \cup M \times \binom{M}{2}$$

What is $f^{-1}(\{x_1, x_2\})$? It's ≤ 1 POINT.

What is $f^{-1}(x_1, \{x_2, x_3\})$? It's ≤ 2 POINTS.

The $d = 1$ Case

Thm: For all $X \subseteq \mathbb{R}^1$ of size n there exists a dist-rainbow subset of size $\Omega(n^{1/3})$.

Proof: Let M be a **MAXIMAL DIST-RAINBOW SET**.

Let $x \in X - M$. WHY IS x NOT IN M !? Either

- ▶ $(\exists x_1, x_2 \in M)[|x - x_1| = |x - x_2|]$, or
- ▶ $(\exists x_1, x_2, x_3 \in M)[|x - x_1| = |x_2 - x_3|]$.

f maps an element of $X - M$ to **reason** $x \notin M$.

$$f : X - M \rightarrow \binom{M}{2} \cup M \times \binom{M}{2}$$

What is $f^{-1}(\{x_1, x_2\})$? It's ≤ 1 POINT.

What is $f^{-1}(x_1, \{x_2, x_3\})$? It's ≤ 2 POINTS.

$$f : X - M \rightarrow \binom{M}{2} \cup M \times \binom{M}{2} \text{ is } \leq 2\text{-to-1.}$$

The $d = 1$ Case- Cont

$f : X - M \rightarrow \binom{M}{2} \cup M \times \binom{M}{2}$ is ≤ 2 -to-1.

The $d = 1$ Case- Cont

$f : X - M \rightarrow \binom{M}{2} \cup M \times \binom{M}{2}$ is ≤ 2 -to-1.

Case 1: $|M| \geq n^{1/3}$ DONE!

The $d = 1$ Case- Cont

$f : X - M \rightarrow \binom{M}{2} \cup M \times \binom{M}{2}$ is ≤ 2 -to-1.

Case 1: $|M| \geq n^{1/3}$ DONE!

Case 2: $|M| \leq n^{1/3}$. So $|X - M| = \Theta(|X|)$. By LEMMA

$$\begin{aligned} |\binom{M}{2} + M \times \binom{M}{2}| &\geq 0.5|X - M| = \Omega(|X|) = \Omega(n) \\ M &\geq \Omega(n^{1/3}) \end{aligned}$$

On Circle

Thm: For all $X \subseteq \mathbb{S}^1$ (the circle) of size n there exists a dist-rainbow subset of size $\Omega(n^{1/3})$.

Proof: Use **MAXIMAL DIST-RAINBOW SET**. Similar Proof.

Better is known

Better is known: In 1975 Komlos, Sulyok, Szemerédi showed:

Thm: For all $X \subseteq \mathbb{S}^1$ or \mathbb{R}^1 of size n there exists a dist-rainbow subset of size $\Omega(n^{1/2})$.

Better is known

Better is known: In 1975 Komlos, Sulyok, Szemerédi showed:

Thm: For all $X \subseteq \mathbb{S}^1$ or \mathbb{R}^1 of size n there exists a dist-rainbow subset of size $\Omega(n^{1/2})$.

This is optimal in \mathbb{S}^1 and \mathbb{R}^1

Better is known

Better is known: In 1975 Komlos, Sulyok, Szemerédi showed:

Thm: For all $X \subseteq \mathbb{S}^1$ or \mathbb{R}^1 of size n there exists a dist-rainbow subset of size $\Omega(n^{1/2})$.

This is optimal in \mathbb{S}^1 and \mathbb{R}^1

Thm: If $X = \{1, \dots, n\}$ then the largest dist-rainbow subset is of size $\leq (1 + o(1))n^{1/2}$.

The $d = 2$ Case

Thm: For all $X \subseteq \mathbb{R}^2$ of size n there exists a dist-rainbow subset of size $\Omega(n^{1/6})$.

Proof: Let M be a **MAXIMAL DIST-RAINBOW SET**.

The $d = 2$ Case

Thm: For all $X \subseteq \mathbb{R}^2$ of size n there exists a dist-rainbow subset of size $\Omega(n^{1/6})$.

Proof: Let M be a **MAXIMAL DIST-RAINBOW SET**.
Let $x \in X - M$. WHY IS x NOT IN M !? Either

The $d = 2$ Case

Thm: For all $X \subseteq \mathbb{R}^2$ of size n there exists a dist-rainbow subset of size $\Omega(n^{1/6})$.

Proof: Let M be a **MAXIMAL DIST-RAINBOW SET**.

Let $x \in X - M$. WHY IS x NOT IN M !? Either

- ▶ $(\exists x_1, x_2 \in M)[|x - x_1| = |x - x_2|]$.

The $d = 2$ Case

Thm: For all $X \subseteq \mathbb{R}^2$ of size n there exists a dist-rainbow subset of size $\Omega(n^{1/6})$.

Proof: Let M be a **MAXIMAL DIST-RAINBOW SET**.

Let $x \in X - M$. WHY IS x NOT IN M !? Either

- ▶ $(\exists x_1, x_2 \in M)[|x - x_1| = |x - x_2|]$.
- ▶ $(\exists x_1, x_2, x_3 \in M)[|x - x_1| = |x_2 - x_3|]$.

f maps an element of $X - M$ to **reason** $x \notin M$.

$$f : X - M \rightarrow \binom{M}{2} \cup M \times \binom{M}{2}$$

What is $f^{-1}(\{x_1, x_2\})$?

The $d = 2$ Case

Thm: For all $X \subseteq \mathbb{R}^2$ of size n there exists a dist-rainbow subset of size $\Omega(n^{1/6})$.

Proof: Let M be a **MAXIMAL DIST-RAINBOW SET**.

Let $x \in X - M$. WHY IS x NOT IN M !? Either

- ▶ $(\exists x_1, x_2 \in M)[|x - x_1| = |x - x_2|]$.
- ▶ $(\exists x_1, x_2, x_3 \in M)[|x - x_1| = |x_2 - x_3|]$.

f maps an element of $X - M$ to **reason** $x \notin M$.

$$f : X - M \rightarrow \binom{M}{2} \cup M \times \binom{M}{2}$$

What is $f^{-1}(\{x_1, x_2\})$? Lies on LINE.

The $d = 2$ Case

Thm: For all $X \subseteq \mathbb{R}^2$ of size n there exists a dist-rainbow subset of size $\Omega(n^{1/6})$.

Proof: Let M be a **MAXIMAL DIST-RAINBOW SET**.

Let $x \in X - M$. WHY IS x NOT IN M !? Either

- ▶ $(\exists x_1, x_2 \in M)[|x - x_1| = |x - x_2|]$.
- ▶ $(\exists x_1, x_2, x_3 \in M)[|x - x_1| = |x_2 - x_3|]$.

f maps an element of $X - M$ to **reason** $x \notin M$.

$$f : X - M \rightarrow \binom{M}{2} \cup M \times \binom{M}{2}$$

What is $f^{-1}(\{x_1, x_2\})$? Lies on LINE.

What is $f^{-1}(x_1, \{x_2, x_3\})$?

The $d = 2$ Case

Thm: For all $X \subseteq \mathbb{R}^2$ of size n there exists a dist-rainbow subset of size $\Omega(n^{1/6})$.

Proof: Let M be a **MAXIMAL DIST-RAINBOW SET**.

Let $x \in X - M$. WHY IS x NOT IN M !? Either

- ▶ $(\exists x_1, x_2 \in M)[|x - x_1| = |x - x_2|]$.
- ▶ $(\exists x_1, x_2, x_3 \in M)[|x - x_1| = |x_2 - x_3|]$.

f maps an element of $X - M$ to **reason** $x \notin M$.

$$f : X - M \rightarrow \binom{M}{2} \cup M \times \binom{M}{2}$$

What is $f^{-1}(\{x_1, x_2\})$? Lies on LINE.

What is $f^{-1}(x_1, \{x_2, x_3\})$? Lies on CIRCLE.

The $d = 2$ Case

Thm: For all $X \subseteq \mathbb{R}^2$ of size n there exists a dist-rainbow subset of size $\Omega(n^{1/6})$.

Proof: Let M be a **MAXIMAL DIST-RAINBOW SET**.

Let $x \in X - M$. WHY IS x NOT IN M !? Either

- ▶ $(\exists x_1, x_2 \in M)[|x - x_1| = |x - x_2|]$.
- ▶ $(\exists x_1, x_2, x_3 \in M)[|x - x_1| = |x_2 - x_3|]$.

f maps an element of $X - M$ to **reason** $x \notin M$.

$$f : X - M \rightarrow \binom{M}{2} \cup M \times \binom{M}{2}$$

What is $f^{-1}(\{x_1, x_2\})$? Lies on LINE.

What is $f^{-1}(x_1, \{x_2, x_3\})$? Lies on CIRCLE.

All INVERSE IMG's lie on LINES or CIRCLES.

The $d = 2$ Case- Cont

$$f : X - M \rightarrow \binom{M}{2} \cup M \times \binom{M}{2}$$

All INVERSE IMG's lie on LINES or CIRCLES. δ TBD.

Cases 1 and 2 induct into line and circle case.

Case 1: $(\exists x_1, x_2)[(f^{-1}(\{x_1, x_2\})| \geq n^\delta]$.

$\geq n^\delta$ points on a line, so rainbow set size $\geq \Omega(n^{\delta/3})$.

The $d = 2$ Case- Cont

$$f : X - M \rightarrow \binom{M}{2} \cup M \times \binom{M}{2}$$

All INVERSE IMG's lie on LINES or CIRCLES. δ TBD.

Cases 1 and 2 induct into line and circle case.

Case 1: $(\exists x_1, x_2)[|f^{-1}(\{x_1, x_2\})| \geq n^\delta]$.

$\geq n^\delta$ points on a line, so rainbow set size $\geq \Omega(n^{\delta/3})$.

Case 2: $(\exists x_1, x_2, x_3)[|f^{-1}(\{x_1, \{x_2, x_3\}\})| \geq n^\delta]$.

$\geq n^\delta$ points on a circle, so rainbow set size $\geq \Omega(n^{\delta/3})$.

The $d = 2$ Case- Cont

$$f : X - M \rightarrow \binom{M}{2} \cup M \times \binom{M}{2}$$

All INVERSE IMG's lie on LINES or CIRCLES. δ TBD.

Cases 1 and 2 induct into line and circle case.

Case 1: $(\exists x_1, x_2)[|f^{-1}(\{x_1, x_2\})| \geq n^\delta]$.

$\geq n^\delta$ points on a line, so rainbow set size $\geq \Omega(n^{\delta/3})$.

Case 2: $(\exists x_1, x_2, x_3)[|f^{-1}(\{x_1, \{x_2, x_3\}\})| \geq n^\delta]$.

$\geq n^\delta$ points on a circle, so rainbow set size $\geq \Omega(n^{\delta/3})$.

Case 3: $|M| \geq n^{1/6}$ DONE!

The $d = 2$ Case- Cont

$$f : X - M \rightarrow \binom{M}{2} \cup M \times \binom{M}{2}$$

All INVERSE IMG's lie on LINES or CIRCLES. δ TBD.

Cases 1 and 2 induct into line and circle case.

Case 1: $(\exists x_1, x_2)[|f^{-1}(\{x_1, x_2\})| \geq n^\delta]$.

$\geq n^\delta$ points on a line, so rainbow set size $\geq \Omega(n^{\delta/3})$.

Case 2: $(\exists x_1, x_2, x_3)[|f^{-1}(\{x_1, \{x_2, x_3\}\})| \geq n^\delta]$.

$\geq n^\delta$ points on a circle, so rainbow set size $\geq \Omega(n^{\delta/3})$.

Case 3: $|M| \geq n^{1/6}$ DONE!

Case 4: Map is $\leq n^\delta$ -to-1 AND $|X - M| = \Theta(|X|)$. By LEMMA

The $d = 2$ Case- Cont

$$f : X - M \rightarrow \binom{M}{2} \cup M \times \binom{M}{2}$$

All INVERSE IMG's lie on LINES or CIRCLES. δ TBD.

Cases 1 and 2 induct into line and circle case.

Case 1: $(\exists x_1, x_2)[|f^{-1}(\{x_1, x_2\})| \geq n^\delta]$.

$\geq n^\delta$ points on a line, so rainbow set size $\geq \Omega(n^{\delta/3})$.

Case 2: $(\exists x_1, x_2, x_3)[|f^{-1}(\{x_1, \{x_2, x_3\}\})| \geq n^\delta]$.

$\geq n^\delta$ points on a circle, so rainbow set size $\geq \Omega(n^{\delta/3})$.

Case 3: $|M| \geq n^{1/6}$ DONE!

Case 4: Map is $\leq n^\delta$ -to-1 AND $|X - M| = \Theta(|X|)$. By LEMMA

$$\begin{aligned} |\binom{M}{2} \cup M \times \binom{M}{2}| &\geq n/n^\delta = n^{1-\delta} \\ |M| &\geq \Omega(n^{(1-\delta)/3}) \end{aligned}$$

The $d = 2$ Case- Cont

$$f : X - M \rightarrow \binom{M}{2} \cup M \times \binom{M}{2}$$

All INVERSE IMG's lie on LINES or CIRCLES. δ TBD.

Cases 1 and 2 induct into line and circle case.

Case 1: $(\exists x_1, x_2)[|f^{-1}(\{x_1, x_2\})| \geq n^\delta]$.

$\geq n^\delta$ points on a line, so rainbow set size $\geq \Omega(n^{\delta/3})$.

Case 2: $(\exists x_1, x_2, x_3)[|f^{-1}(\{x_1, \{x_2, x_3\}\})| \geq n^\delta]$.

$\geq n^\delta$ points on a circle, so rainbow set size $\geq \Omega(n^{\delta/3})$.

Case 3: $|M| \geq n^{1/6}$ DONE!

Case 4: Map is $\leq n^\delta$ -to-1 AND $|X - M| = \Theta(|X|)$. By LEMMA

$$\begin{aligned} |\binom{M}{2} \cup M \times \binom{M}{2}| &\geq n/n^\delta = n^{1-\delta} \\ |M| &\geq \Omega(n^{(1-\delta)/3}) \end{aligned}$$

Set $\delta/3 = (1 - \delta)/3$. $\delta = 1/2$. Get $\Omega(n^{1/6})$.

On Sphere

Thm: For all $X \subseteq \mathbb{S}^2$ (surface of sphere) of size n there exists a dist-rainbow subset of size $\Omega(n^{1/6})$.

Proof: Use **MAXIMAL DIST-RAINBOW SET**. Similar Proof.

On Sphere

Thm: For all $X \subseteq \mathbb{S}^2$ (surface of sphere) of size n there exists a dist-rainbow subset of size $\Omega(n^{1/6})$.

Proof: Use **MAXIMAL DIST-RAINBOW SET**. Similar Proof.

Note: Better is known: Charalambides showed $\Omega(n^{1/3})$.

General d Case

Thm:

For all $X \subseteq \mathbb{R}^d$ of size $n \exists$ dist-rainbow subset of size $\Omega(n^{1/3d})$.

For all $X \subseteq \mathbb{S}^d$ of size $n \exists$ dist-rainbow subset of size $\Omega(n^{1/3d})$.

General d Case

Thm:

For all $X \subseteq \mathbb{R}^d$ of size $n \exists$ dist-rainbow subset of size $\Omega(n^{1/3d})$.

For all $X \subseteq \mathbb{S}^d$ of size $n \exists$ dist-rainbow subset of size $\Omega(n^{1/3d})$.

Proof: Use **MAXIMAL DIST-RAINBOW SET** and induction.
Need result on \mathbb{S}^d and \mathbb{R}^d to get result for \mathbb{S}^{d+1} and \mathbb{R}^{d+1} .

General d Case

Thm:

For all $X \subseteq \mathbb{R}^d$ of size $n \ni$ dist-rainbow subset of size $\Omega(n^{1/3d})$.

For all $X \subseteq \mathbb{S}^d$ of size $n \ni$ dist-rainbow subset of size $\Omega(n^{1/3d})$.

Proof: Use **MAXIMAL DIST-RAINBOW SET** and induction.
Need result on \mathbb{S}^d and \mathbb{R}^d to get result for \mathbb{S}^{d+1} and \mathbb{R}^{d+1} .

Note: Better is known. In 1995 Thiele showed $\Omega(n^{1/(3d-2)})$. But WE improved that!

General d Case- Much Better

Thm: For all $d \geq 2$, for all $X \subseteq \mathbb{R}^d$ of size n there exists a dist-rainbow subset of size $\Omega(n^{1/(3d-3)}(\log n)^{\frac{1}{3} - \frac{2}{3d-3}})$.

General d Case- Much Better

Thm: For all $d \geq 2$, for all $X \subseteq \mathbb{R}^d$ of size n there exists a dist-rainbow subset of size $\Omega(n^{1/(3d-3)}(\log n)^{\frac{1}{3}-\frac{2}{3d-3}})$.

Proof: Use **VARIANT ON MAX DIST-RAINBOW SET**

d	$n^{1/3d}$	$n^{1/(3d-3)}(\log n)^{\frac{1}{3}-\frac{2}{3d-3}}$
1	$n^{1/3}$	--
2	$n^{1/6}$	$n^{1/3}(\log n)^{-1/3}$
3	$n^{1/9}$	$n^{1/6}(\log n)^0$
4	$n^{1/12}$	$n^{1/9}(\log n)^{1/12}$
5	$n^{1/15}$	$n^{1/12}(\log n)^{1/6}$
6	$n^{1/18}$	$n^{1/15}(\log n)^{1/5}$

General d Case- Much Better

Thm: For all $d \geq 2$, for all $X \subseteq \mathbb{R}^d$ of size n there exists a dist-rainbow subset of size $\Omega(n^{1/(3d-3)}(\log n)^{\frac{1}{3}-\frac{2}{3d-3}})$.

Proof: Use **VARIANT ON MAX DIST-RAINBOW SET**

d	$n^{1/3d}$	$n^{1/(3d-3)}(\log n)^{\frac{1}{3}-\frac{2}{3d-3}}$
1	$n^{1/3}$	--
2	$n^{1/6}$	$n^{1/3}(\log n)^{-1/3}$
3	$n^{1/9}$	$n^{1/6}(\log n)^0$
4	$n^{1/12}$	$n^{1/9}(\log n)^{1/12}$
5	$n^{1/15}$	$n^{1/12}(\log n)^{1/6}$
6	$n^{1/18}$	$n^{1/15}(\log n)^{1/5}$

Can we do better? Best we can hope for is roughly $n^{1/d}$.

Area- $d = 2$ Case

Thm: For all $X \subseteq \mathbb{R}^2$ of size n , no three colinear, \exists area-rainbow set of size $\Omega(n^{1/5})$.

Area- $d = 2$ Case

Thm: For all $X \subseteq \mathbb{R}^2$ of size n , no three colinear, \exists area-rainbow set of size $\Omega(n^{1/5})$.

Proof: Let M be a **MAXIMAL AREA-RAINBOW SET**.

Area- $d = 2$ Case

Thm: For all $X \subseteq \mathbb{R}^2$ of size n , no three colinear, \exists area-rainbow set of size $\Omega(n^{1/5})$.

Proof: Let M be a **MAXIMAL AREA-RAINBOW SET**.

Let $x \in X - M$. WHY IS x NOT IN M !? Either

Area- $d = 2$ Case

Thm: For all $X \subseteq \mathbb{R}^2$ of size n , no three colinear, \exists area-rainbow set of size $\Omega(n^{1/5})$.

Proof: Let M be a **MAXIMAL AREA-RAINBOW SET**.

Let $x \in X - M$. WHY IS x NOT IN M !? Either

- ▶ $(\exists x_1, x_2, x_3 \in M)[\text{AREA}(x, x_1, x_2) = \text{AREA}(x, x_1, x_3)]$.

Area- $d = 2$ Case

Thm: For all $X \subseteq \mathbb{R}^2$ of size n , no three colinear, \exists area-rainbow set of size $\Omega(n^{1/5})$.

Proof: Let M be a **MAXIMAL AREA-RAINBOW SET**.

Let $x \in X - M$. WHY IS x NOT IN M !? Either

- ▶ $(\exists x_1, x_2, x_3 \in M)[\text{AREA}(x, x_1, x_2) = \text{AREA}(x, x_1, x_3)]$.
- ▶ $(\exists x_1, x_2, x_3, x_4 \in M)[\text{AREA}(x, x_1, x_2) = \text{AREA}(x, x_3, x_4)]$.

Area- $d = 2$ Case

Thm: For all $X \subseteq \mathbb{R}^2$ of size n , no three colinear, \exists area-rainbow set of size $\Omega(n^{1/5})$.

Proof: Let M be a **MAXIMAL AREA-RAINBOW SET**.

Let $x \in X - M$. WHY IS x NOT IN M !? Either

- ▶ $(\exists x_1, x_2, x_3 \in M)[\text{AREA}(x, x_1, x_2) = \text{AREA}(x, x_1, x_3)]$.
- ▶ $(\exists x_1, x_2, x_3, x_4 \in M)[\text{AREA}(x, x_1, x_2) = \text{AREA}(x, x_3, x_4)]$.
- ▶ $(\exists x_1, x_2, x_3, x_4, x_5 \in M)[\text{AREA}(x, x_1, x_2) = \text{AREA}(x_3, x_4, x_5)]$.

Area- $d = 2$ Case

Thm: For all $X \subseteq \mathbb{R}^2$ of size n , no three colinear, \exists area-rainbow set of size $\Omega(n^{1/5})$.

Proof: Let M be a **MAXIMAL AREA-RAINBOW SET**.

Let $x \in X - M$. WHY IS x NOT IN M !? Either

- ▶ $(\exists x_1, x_2, x_3 \in M)[\text{AREA}(x, x_1, x_2) = \text{AREA}(x, x_1, x_3)]$.
- ▶ $(\exists x_1, x_2, x_3, x_4 \in M)[\text{AREA}(x, x_1, x_2) = \text{AREA}(x, x_3, x_4)]$.
- ▶ $(\exists x_1, x_2, x_3, x_4, x_5 \in M)[\text{AREA}(x, x_1, x_2) = \text{AREA}(x_3, x_4, x_5)]$.

f maps an element of $X - M$ to **reason** $x \notin M$.

Area- $d = 2$ Case

Thm: For all $X \subseteq \mathbb{R}^2$ of size n , no three colinear, \exists area-rainbow set of size $\Omega(n^{1/5})$.

Proof: Let M be a **MAXIMAL AREA-RAINBOW SET**.

Let $x \in X - M$. WHY IS x NOT IN M !? Either

- ▶ $(\exists x_1, x_2, x_3 \in M)[\text{AREA}(x, x_1, x_2) = \text{AREA}(x, x_1, x_3)]$.
- ▶ $(\exists x_1, x_2, x_3, x_4 \in M)[\text{AREA}(x, x_1, x_2) = \text{AREA}(x, x_3, x_4)]$.
- ▶ $(\exists x_1, x_2, x_3, x_4, x_5 \in M)[\text{AREA}(x, x_1, x_2) = \text{AREA}(x_3, x_4, x_5)]$.

f maps an element of $X - M$ to **reason** $x \notin M$.

$f : X - M \rightarrow \binom{M}{2} \times \binom{M}{2} \cup \binom{M}{2} \times \binom{M}{3}$. Recall that

Area- $d = 2$ Case

Thm: For all $X \subseteq \mathbb{R}^2$ of size n , no three colinear, \exists area-rainbow set of size $\Omega(n^{1/5})$.

Proof: Let M be a **MAXIMAL AREA-RAINBOW SET**.

Let $x \in X - M$. WHY IS x NOT IN M !? Either

- ▶ $(\exists x_1, x_2, x_3 \in M)[\text{AREA}(x, x_1, x_2) = \text{AREA}(x, x_1, x_3)]$.
- ▶ $(\exists x_1, x_2, x_3, x_4 \in M)[\text{AREA}(x, x_1, x_2) = \text{AREA}(x, x_3, x_4)]$.
- ▶ $(\exists x_1, x_2, x_3, x_4, x_5 \in M)[\text{AREA}(x, x_1, x_2) = \text{AREA}(x_3, x_4, x_5)]$.

f maps an element of $X - M$ to **reason** $x \notin M$.

$f : X - M \rightarrow \binom{M}{2} \times \binom{M}{2} \cup \binom{M}{2} \times \binom{M}{3}$. Recall that

What is $f^{-1}(\{x_1, x_2\}, \{x_1, x_3\})$? SEE NEXT SLIDE FOR GEOM LEMMA.

Lemma On Area

Lemma: Let L_1 and L_2 be lines in \mathbb{R}^2 .

$$\{p : \text{AREA}(L_1, p) = \text{AREA}(L_2, p)\}$$

is a line.

Lemma On Area

Lemma: Let L_1 and L_2 be lines in \mathbb{R}^2 .

$$\{p : \text{AREA}(L_1, p) = \text{AREA}(L_2, p)\}$$

is a line.

Sketch: $\text{AREA}(L_1, p) = \text{AREA}(L_2, p)$ iff

$|L_1| \times |L_1 - p| = |L_2| \times |L_2 - p|$ iff $\frac{|L_1 - p|}{|L_2 - p|} = \frac{|L_1|}{|L_2|}$. This is a line.

(Reboot) Area- $d = 2$ Case

Thm: For all $X \subseteq \mathbb{R}^2$ of size n , no three colinear, \exists area-rainbow set of size $\Omega(n^{1/5})$.

(Reboot) Area- $d = 2$ Case

Thm: For all $X \subseteq \mathbb{R}^2$ of size n , no three colinear, \exists area-rainbow set of size $\Omega(n^{1/5})$.

Proof: Let M be a **MAXIMAL AREA-RAINBOW SET**.

(Reboot) Area- $d = 2$ Case

Thm: For all $X \subseteq \mathbb{R}^2$ of size n , no three colinear, \exists area-rainbow set of size $\Omega(n^{1/5})$.

Proof: Let M be a **MAXIMAL AREA-RAINBOW SET**.

Let $x \in X - M$. WHY IS x NOT IN M !? Either

(Reboot) Area- $d = 2$ Case

Thm: For all $X \subseteq \mathbb{R}^2$ of size n , no three colinear, \exists area-rainbow set of size $\Omega(n^{1/5})$.

Proof: Let M be a **MAXIMAL AREA-RAINBOW SET**.

Let $x \in X - M$. WHY IS x NOT IN M !? Either

- ▶ $(\exists x_1, x_2, x_3 \in M)[\text{AREA}(x, x_1, x_2) = \text{AREA}(x, x_1, x_3)]$.

(Reboot) Area- $d = 2$ Case

Thm: For all $X \subseteq \mathbb{R}^2$ of size n , no three colinear, \exists area-rainbow set of size $\Omega(n^{1/5})$.

Proof: Let M be a **MAXIMAL AREA-RAINBOW SET**.

Let $x \in X - M$. WHY IS x NOT IN M !? Either

- ▶ $(\exists x_1, x_2, x_3 \in M)[\text{AREA}(x, x_1, x_2) = \text{AREA}(x, x_1, x_3)]$.
- ▶ $(\exists x_1, x_2, x_3, x_4 \in M)[\text{AREA}(x, x_1, x_2) = \text{AREA}(x, x_3, x_4)]$.

(Reboot) Area- $d = 2$ Case

Thm: For all $X \subseteq \mathbb{R}^2$ of size n , no three colinear, \exists area-rainbow set of size $\Omega(n^{1/5})$.

Proof: Let M be a **MAXIMAL AREA-RAINBOW SET**.

Let $x \in X - M$. WHY IS x NOT IN M !? Either

- ▶ $(\exists x_1, x_2, x_3 \in M)[\text{AREA}(x, x_1, x_2) = \text{AREA}(x, x_1, x_3)]$.
- ▶ $(\exists x_1, x_2, x_3, x_4 \in M)[\text{AREA}(x, x_1, x_2) = \text{AREA}(x, x_3, x_4)]$.
- ▶ $(\exists x_1, x_2, x_3, x_4, x_5 \in M)[\text{AREA}(x, x_1, x_2) = \text{AREA}(x_3, x_4, x_5)]$.

(Reboot) Area- $d = 2$ Case

Thm: For all $X \subseteq \mathbb{R}^2$ of size n , no three colinear, \exists area-rainbow set of size $\Omega(n^{1/5})$.

Proof: Let M be a **MAXIMAL AREA-RAINBOW SET**.

Let $x \in X - M$. WHY IS x NOT IN M !? Either

- ▶ $(\exists x_1, x_2, x_3 \in M)[\text{AREA}(x, x_1, x_2) = \text{AREA}(x, x_1, x_3)]$.
- ▶ $(\exists x_1, x_2, x_3, x_4 \in M)[\text{AREA}(x, x_1, x_2) = \text{AREA}(x, x_3, x_4)]$.
- ▶ $(\exists x_1, x_2, x_3, x_4, x_5 \in M)[\text{AREA}(x, x_1, x_2) = \text{AREA}(x_3, x_4, x_5)]$.

f maps an element of $X - M$ to **reason** $x \notin M$.

(Reboot) Area- $d = 2$ Case

Thm: For all $X \subseteq \mathbb{R}^2$ of size n , no three colinear, \exists area-rainbow set of size $\Omega(n^{1/5})$.

Proof: Let M be a **MAXIMAL AREA-RAINBOW SET**.

Let $x \in X - M$. WHY IS x NOT IN M !? Either

- ▶ $(\exists x_1, x_2, x_3 \in M)[\text{AREA}(x, x_1, x_2) = \text{AREA}(x, x_1, x_3)]$.
- ▶ $(\exists x_1, x_2, x_3, x_4 \in M)[\text{AREA}(x, x_1, x_2) = \text{AREA}(x, x_3, x_4)]$.
- ▶ $(\exists x_1, x_2, x_3, x_4, x_5 \in M)[\text{AREA}(x, x_1, x_2) = \text{AREA}(x_3, x_4, x_5)]$.

f maps an element of $X - M$ to **reason** $x \notin M$.

$f : X - M \rightarrow \binom{M}{2} \times \binom{M}{2} \cup \binom{M}{2} \times \binom{M}{3}$. Recall that

(Reboot) Area- $d = 2$ Case

Thm: For all $X \subseteq \mathbb{R}^2$ of size n , no three colinear, \exists area-rainbow set of size $\Omega(n^{1/5})$.

Proof: Let M be a **MAXIMAL AREA-RAINBOW SET**.

Let $x \in X - M$. WHY IS x NOT IN M !? Either

- ▶ $(\exists x_1, x_2, x_3 \in M)[\text{AREA}(x, x_1, x_2) = \text{AREA}(x, x_1, x_3)]$.
- ▶ $(\exists x_1, x_2, x_3, x_4 \in M)[\text{AREA}(x, x_1, x_2) = \text{AREA}(x, x_3, x_4)]$.
- ▶ $(\exists x_1, x_2, x_3, x_4, x_5 \in M)[\text{AREA}(x, x_1, x_2) = \text{AREA}(x_3, x_4, x_5)]$.

f maps an element of $X - M$ to **reason** $x \notin M$.

$f : X - M \rightarrow \binom{M}{2} \times \binom{M}{2} \cup \binom{M}{2} \times \binom{M}{3}$. Recall that
What is $f^{-1}(\{x_1, x_2\}, \{x_1, x_3\})$?

(Reboot) Area- $d = 2$ Case

Thm: For all $X \subseteq \mathbb{R}^2$ of size n , no three colinear, \exists area-rainbow set of size $\Omega(n^{1/5})$.

Proof: Let M be a **MAXIMAL AREA-RAINBOW SET**.

Let $x \in X - M$. WHY IS x NOT IN M !? Either

- ▶ $(\exists x_1, x_2, x_3 \in M)[\text{AREA}(x, x_1, x_2) = \text{AREA}(x, x_1, x_3)]$.
- ▶ $(\exists x_1, x_2, x_3, x_4 \in M)[\text{AREA}(x, x_1, x_2) = \text{AREA}(x, x_3, x_4)]$.
- ▶ $(\exists x_1, x_2, x_3, x_4, x_5 \in M)[\text{AREA}(x, x_1, x_2) = \text{AREA}(x_3, x_4, x_5)]$.

f maps an element of $X - M$ to **reason** $x \notin M$.

$f : X - M \rightarrow \binom{M}{2} \times \binom{M}{2} \cup \binom{M}{2} \times \binom{M}{3}$. Recall that

What is $f^{-1}(\{x_1, x_2\}, \{x_1, x_3\})$? By Lemma all points on it are on a line- so ≤ 2 points. FINITE.

(Reboot) Area- $d = 2$ Case

Thm: For all $X \subseteq \mathbb{R}^2$ of size n , no three colinear, \exists area-rainbow set of size $\Omega(n^{1/5})$.

Proof: Let M be a **MAXIMAL AREA-RAINBOW SET**.

Let $x \in X - M$. WHY IS x NOT IN M !? Either

- ▶ $(\exists x_1, x_2, x_3 \in M)[\text{AREA}(x, x_1, x_2) = \text{AREA}(x, x_1, x_3)]$.
- ▶ $(\exists x_1, x_2, x_3, x_4 \in M)[\text{AREA}(x, x_1, x_2) = \text{AREA}(x, x_3, x_4)]$.
- ▶ $(\exists x_1, x_2, x_3, x_4, x_5 \in M)[\text{AREA}(x, x_1, x_2) = \text{AREA}(x_3, x_4, x_5)]$.

f maps an element of $X - M$ to **reason** $x \notin M$.

$f : X - M \rightarrow \binom{M}{2} \times \binom{M}{2} \cup \binom{M}{2} \times \binom{M}{3}$. Recall that

What is $f^{-1}(\{x_1, x_2\}, \{x_1, x_3\})$? By Lemma all points on it are on a line- so ≤ 2 points. FINITE.

What is $f^{-1}(\{x_1, x_2\}, \{x_3, x_4\})$?

(Reboot) Area- $d = 2$ Case

Thm: For all $X \subseteq \mathbb{R}^2$ of size n , no three colinear, \exists area-rainbow set of size $\Omega(n^{1/5})$.

Proof: Let M be a **MAXIMAL AREA-RAINBOW SET**.

Let $x \in X - M$. WHY IS x NOT IN M !? Either

- ▶ $(\exists x_1, x_2, x_3 \in M)[\text{AREA}(x, x_1, x_2) = \text{AREA}(x, x_1, x_3)]$.
- ▶ $(\exists x_1, x_2, x_3, x_4 \in M)[\text{AREA}(x, x_1, x_2) = \text{AREA}(x, x_3, x_4)]$.
- ▶ $(\exists x_1, x_2, x_3, x_4, x_5 \in M)[\text{AREA}(x, x_1, x_2) = \text{AREA}(x_3, x_4, x_5)]$.

f maps an element of $X - M$ to **reason** $x \notin M$.

$f : X - M \rightarrow \binom{M}{2} \times \binom{M}{2} \cup \binom{M}{2} \times \binom{M}{3}$. Recall that

What is $f^{-1}(\{x_1, x_2\}, \{x_1, x_3\})$? By Lemma all points on it are on a line- so ≤ 2 points. FINITE.

What is $f^{-1}(\{x_1, x_2\}, \{x_3, x_4\})$? By Lemma all points on it are on a line- so ≤ 2 points. FINITE.

(Reboot) Area- $d = 2$ Case

Thm: For all $X \subseteq \mathbb{R}^2$ of size n , no three colinear, \exists area-rainbow set of size $\Omega(n^{1/5})$.

Proof: Let M be a **MAXIMAL AREA-RAINBOW SET**.

Let $x \in X - M$. WHY IS x NOT IN M !? Either

- ▶ $(\exists x_1, x_2, x_3 \in M)[\text{AREA}(x, x_1, x_2) = \text{AREA}(x, x_1, x_3)]$.
- ▶ $(\exists x_1, x_2, x_3, x_4 \in M)[\text{AREA}(x, x_1, x_2) = \text{AREA}(x, x_3, x_4)]$.
- ▶ $(\exists x_1, x_2, x_3, x_4, x_5 \in M)[\text{AREA}(x, x_1, x_2) = \text{AREA}(x_3, x_4, x_5)]$.

f maps an element of $X - M$ to **reason** $x \notin M$.

$f : X - M \rightarrow \binom{M}{2} \times \binom{M}{2} \cup \binom{M}{2} \times \binom{M}{3}$. Recall that

What is $f^{-1}(\{x_1, x_2\}, \{x_1, x_3\})$? By Lemma all points on it are on a line- so ≤ 2 points. FINITE.

What is $f^{-1}(\{x_1, x_2\}, \{x_3, x_4\})$? By Lemma all points on it are on a line- so ≤ 2 points. FINITE.

What is $f^{-1}(\{x_1, x_2\}, \{x_3, x_4, x_5\})$?

(Reboot) Area- $d = 2$ Case

Thm: For all $X \subseteq \mathbb{R}^2$ of size n , no three colinear, \exists area-rainbow set of size $\Omega(n^{1/5})$.

Proof: Let M be a **MAXIMAL AREA-RAINBOW SET**.

Let $x \in X - M$. WHY IS x NOT IN M !? Either

- ▶ $(\exists x_1, x_2, x_3 \in M)[\text{AREA}(x, x_1, x_2) = \text{AREA}(x, x_1, x_3)]$.
- ▶ $(\exists x_1, x_2, x_3, x_4 \in M)[\text{AREA}(x, x_1, x_2) = \text{AREA}(x, x_3, x_4)]$.
- ▶ $(\exists x_1, x_2, x_3, x_4, x_5 \in M)[\text{AREA}(x, x_1, x_2) = \text{AREA}(x_3, x_4, x_5)]$.


f maps an element of $X - M$ to **reason** $x \notin M$.

$f : X - M \rightarrow \binom{M}{2} \times \binom{M}{2} \cup \binom{M}{2} \times \binom{M}{3}$. Recall that

What is $f^{-1}(\{x_1, x_2\}, \{x_1, x_3\})$? By Lemma all points on it are on a line- so ≤ 2 points. FINITE.

What is $f^{-1}(\{x_1, x_2\}, \{x_3, x_4\})$? By Lemma all points on it are on a line- so ≤ 2 points. FINITE.

What is $f^{-1}(\{x_1, x_2\}, \{x_3, x_4, x_5\})$? By Lemma all points on it are on a line- so ≤ 2 points. FINITE.

$f : X - M \rightarrow \binom{M}{2} \times \binom{M}{2} \cup \binom{M}{2} \times \binom{M}{3}$ FINITE-to-1. 

Area $d = 2$ Case- Cont

$f : X - M \rightarrow \binom{M}{2} \times \binom{M}{2} \cup \binom{M}{2} \times \binom{M}{3}$ is FINITE-to-1.

Case 1: $|M| \geq n^{1/5}$ DONE!

Case 2: $|M| \leq n^{1/5}$. Then $|X - M| = \Theta(|X|)$. Since MAP is finite-to-1, by LEMMA

$$\begin{aligned} |\binom{M}{2} \times \binom{M}{2} \cup \binom{M}{2} \times \binom{M}{3}| &\geq \Omega(|X - M|) = \Omega(|X|) = \Omega(n) \\ |M| &\geq \Omega(n^{1/5}) \end{aligned}$$

Volume $d = 3$

Thm: For all $X \subseteq \mathbb{R}^3$ of size n , no four on a plane, there exists Vol-rainbow set of size $\Omega(n^\delta)$. (δ TBD)
Similar. Left for the reader.

KEY to These Proofs

KEY to These Proofs

1. Used **MAXIMAL a -RAINBOW SET** M .

KEY to These Proofs

1. Used **MAXIMAL a -RAINBOW SET** M .
2. Used Map f from $x \in X - M$ to the reason x is NOT in M .

KEY to These Proofs

1. Used **MAXIMAL a -RAINBOW SET** M .
2. Used Map f from $x \in X - M$ to the reason x is NOT in M .
3. Looked at **INVERSE IMAGES** of that map.

KEY to These Proofs

1. Used **MAXIMAL a -RAINBOW SET** M .
2. Used Map f from $x \in X - M$ to the reason x is NOT in M .
3. Looked at **INVERSE IMAGES** of that map.
4. Either:
All INVERSE IMG's are small, so use LEMMA.
OR
Some INVERSE IMG's are large subsets of \mathbb{R}^d or \mathbb{S}^d , so induct.

Area- $d = 3$ Case

Thm: For all $X \subseteq \mathbb{R}^3$ of size n , no three colinear, there exists Area-rainbow set of size $\Omega(n^\delta)$. (δ TBD)

Area- $d = 3$ Case

Thm: For all $X \subseteq \mathbb{R}^3$ of size n , no three colinear, there exists Area-rainbow set of size $\Omega(n^\delta)$. (δ TBD)

Proof: Let M be a **MAXIMAL AREA-RAINBOW SET**.

Area- $d = 3$ Case

Thm: For all $X \subseteq \mathbb{R}^3$ of size n , no three colinear, there exists Area-rainbow set of size $\Omega(n^\delta)$. (δ TBD)

Proof: Let M be a **MAXIMAL AREA-RAINBOW SET**.

Let $x \in X - M$. WHY IS x NOT IN M !? Either

Area- $d = 3$ Case

Thm: For all $X \subseteq \mathbb{R}^3$ of size n , no three colinear, there exists Area-rainbow set of size $\Omega(n^\delta)$. (δ TBD)

Proof: Let M be a **MAXIMAL AREA-RAINBOW SET**.

Let $x \in X - M$. WHY IS x NOT IN M !? Either

- ▶ $(\exists x_1, x_2, x_3 \in M)[\text{AREA}(x, x_1, x_2) = \text{AREA}(x, x_1, x_3)]$.

Area- $d = 3$ Case

Thm: For all $X \subseteq \mathbb{R}^3$ of size n , no three colinear, there exists Area-rainbow set of size $\Omega(n^\delta)$. (δ TBD)

Proof: Let M be a **MAXIMAL AREA-RAINBOW SET**.

Let $x \in X - M$. WHY IS x NOT IN M !? Either

- ▶ $(\exists x_1, x_2, x_3 \in M)[\text{AREA}(x, x_1, x_2) = \text{AREA}(x, x_1, x_3)]$.
- ▶ $(\exists x_1, x_2, x_3, x_4 \in M)[\text{AREA}(x, x_1, x_2) = \text{AREA}(x, x_3, x_4)]$.

Area- $d = 3$ Case

Thm: For all $X \subseteq \mathbb{R}^3$ of size n , no three colinear, there exists Area-rainbow set of size $\Omega(n^\delta)$. (δ TBD)

Proof: Let M be a **MAXIMAL AREA-RAINBOW SET**.

Let $x \in X - M$. WHY IS x NOT IN M !? Either

- ▶ $(\exists x_1, x_2, x_3 \in M)[\text{AREA}(x, x_1, x_2) = \text{AREA}(x, x_1, x_3)]$.
- ▶ $(\exists x_1, x_2, x_3, x_4 \in M)[\text{AREA}(x, x_1, x_2) = \text{AREA}(x, x_3, x_4)]$.
- ▶ $(\exists x_1, x_2, x_3, x_4, x_5 \in M)[\text{AREA}(x, x_1, x_2) = \text{AREA}(x_3, x_4, x_5)]$.

Area- $d = 3$ Case

Thm: For all $X \subseteq \mathbb{R}^3$ of size n , no three colinear, there exists Area-rainbow set of size $\Omega(n^\delta)$. (δ TBD)

Proof: Let M be a **MAXIMAL AREA-RAINBOW SET**.

Let $x \in X - M$. WHY IS x NOT IN M !? Either

- ▶ $(\exists x_1, x_2, x_3 \in M)[\text{AREA}(x, x_1, x_2) = \text{AREA}(x, x_1, x_3)]$.
- ▶ $(\exists x_1, x_2, x_3, x_4 \in M)[\text{AREA}(x, x_1, x_2) = \text{AREA}(x, x_3, x_4)]$.
- ▶ $(\exists x_1, x_2, x_3, x_4, x_5 \in M)[\text{AREA}(x, x_1, x_2) = \text{AREA}(x_3, x_4, x_5)]$.

f maps an element of $X - M$ to **reason** $x \notin M$.

Area- $d = 3$ Case

Thm: For all $X \subseteq \mathbb{R}^3$ of size n , no three colinear, there exists Area-rainbow set of size $\Omega(n^\delta)$. (δ TBD)

Proof: Let M be a **MAXIMAL AREA-RAINBOW SET**.

Let $x \in X - M$. WHY IS x NOT IN M !? Either

- ▶ $(\exists x_1, x_2, x_3 \in M)[\text{AREA}(x, x_1, x_2) = \text{AREA}(x, x_1, x_3)]$.
- ▶ $(\exists x_1, x_2, x_3, x_4 \in M)[\text{AREA}(x, x_1, x_2) = \text{AREA}(x, x_3, x_4)]$.
- ▶ $(\exists x_1, x_2, x_3, x_4, x_5 \in M)[\text{AREA}(x, x_1, x_2) = \text{AREA}(x_3, x_4, x_5)]$.

f maps an element of $X - M$ to **reason** $x \notin M$.

$$f : X - M \rightarrow \binom{M}{2} \times \binom{M}{2} \cup \binom{M}{2} \times \binom{M}{3}.$$

Area- $d = 3$ Case

Thm: For all $X \subseteq \mathbb{R}^3$ of size n , no three colinear, there exists Area-rainbow set of size $\Omega(n^\delta)$. (δ TBD)

Proof: Let M be a **MAXIMAL AREA-RAINBOW SET**.

Let $x \in X - M$. WHY IS x NOT IN M !? Either

- ▶ $(\exists x_1, x_2, x_3 \in M)[\text{AREA}(x, x_1, x_2) = \text{AREA}(x, x_1, x_3)]$.
- ▶ $(\exists x_1, x_2, x_3, x_4 \in M)[\text{AREA}(x, x_1, x_2) = \text{AREA}(x, x_3, x_4)]$.
- ▶ $(\exists x_1, x_2, x_3, x_4, x_5 \in M)[\text{AREA}(x, x_1, x_2) = \text{AREA}(x_3, x_4, x_5)]$.

f maps an element of $X - M$ to **reason** $x \notin M$.

$$f : X - M \rightarrow \binom{M}{2} \times \binom{M}{2} \cup \binom{M}{2} \times \binom{M}{3}.$$

What is $f^{-1}(\{\{x_1, x_2\}, \{x_1, x_3\}\})$?

Area- $d = 3$ Case

Thm: For all $X \subseteq \mathbb{R}^3$ of size n , no three colinear, there exists Area-rainbow set of size $\Omega(n^\delta)$. (δ TBD)

Proof: Let M be a **MAXIMAL AREA-RAINBOW SET**.

Let $x \in X - M$. WHY IS x NOT IN M !? Either

- ▶ $(\exists x_1, x_2, x_3 \in M)[\text{AREA}(x, x_1, x_2) = \text{AREA}(x, x_1, x_3)]$.
- ▶ $(\exists x_1, x_2, x_3, x_4 \in M)[\text{AREA}(x, x_1, x_2) = \text{AREA}(x, x_3, x_4)]$.
- ▶ $(\exists x_1, x_2, x_3, x_4, x_5 \in M)[\text{AREA}(x, x_1, x_2) = \text{AREA}(x_3, x_4, x_5)]$.

f maps an element of $X - M$ to **reason** $x \notin M$.

$f : X - M \rightarrow \binom{M}{2} \times \binom{M}{2} \cup \binom{M}{2} \times \binom{M}{3}$.

What is $f^{-1}(\{\{x_1, x_2\}, \{x_1, x_3\}\})$? THIS IS HARD!

Area- $d = 3$ Case

Thm: For all $X \subseteq \mathbb{R}^3$ of size n , no three colinear, there exists Area-rainbow set of size $\Omega(n^\delta)$. (δ TBD)

Proof: Let M be a **MAXIMAL AREA-RAINBOW SET**.

Let $x \in X - M$. WHY IS x NOT IN M !? Either

- ▶ $(\exists x_1, x_2, x_3 \in M)[\text{AREA}(x, x_1, x_2) = \text{AREA}(x, x_1, x_3)]$.
- ▶ $(\exists x_1, x_2, x_3, x_4 \in M)[\text{AREA}(x, x_1, x_2) = \text{AREA}(x, x_3, x_4)]$.
- ▶ $(\exists x_1, x_2, x_3, x_4, x_5 \in M)[\text{AREA}(x, x_1, x_2) = \text{AREA}(x_3, x_4, x_5)]$.

f maps an element of $X - M$ to **reason** $x \notin M$.

$f : X - M \rightarrow \binom{M}{2} \times \binom{M}{2} \cup \binom{M}{2} \times \binom{M}{3}$.

What is $f^{-1}(\{\{x_1, x_2\}, \{x_1, x_3\}\})$? THIS IS HARD!

What is $f^{-1}(\{x_1, x_2\}, \{x_3, x_4\})$?

Area- $d = 3$ Case

Thm: For all $X \subseteq \mathbb{R}^3$ of size n , no three colinear, there exists Area-rainbow set of size $\Omega(n^\delta)$. (δ TBD)

Proof: Let M be a **MAXIMAL AREA-RAINBOW SET**.

Let $x \in X - M$. WHY IS x NOT IN M !? Either

- ▶ $(\exists x_1, x_2, x_3 \in M)[\text{AREA}(x, x_1, x_2) = \text{AREA}(x, x_1, x_3)]$.
- ▶ $(\exists x_1, x_2, x_3, x_4 \in M)[\text{AREA}(x, x_1, x_2) = \text{AREA}(x, x_3, x_4)]$.
- ▶ $(\exists x_1, x_2, x_3, x_4, x_5 \in M)[\text{AREA}(x, x_1, x_2) = \text{AREA}(x_3, x_4, x_5)]$.

f maps an element of $X - M$ to **reason** $x \notin M$.

$f : X - M \rightarrow \binom{M}{2} \times \binom{M}{2} \cup \binom{M}{2} \times \binom{M}{3}$.

What is $f^{-1}(\{\{x_1, x_2\}, \{x_1, x_3\}\})$? THIS IS HARD!

What is $f^{-1}(\{x_1, x_2\}, \{x_3, x_4\})$? THIS IS HARD!

Area- $d = 3$ Case

Thm: For all $X \subseteq \mathbb{R}^3$ of size n , no three colinear, there exists Area-rainbow set of size $\Omega(n^\delta)$. (δ TBD)

Proof: Let M be a **MAXIMAL AREA-RAINBOW SET**.

Let $x \in X - M$. WHY IS x NOT IN M !? Either

- ▶ $(\exists x_1, x_2, x_3 \in M)[\text{AREA}(x, x_1, x_2) = \text{AREA}(x, x_1, x_3)]$.
- ▶ $(\exists x_1, x_2, x_3, x_4 \in M)[\text{AREA}(x, x_1, x_2) = \text{AREA}(x, x_3, x_4)]$.
- ▶ $(\exists x_1, x_2, x_3, x_4, x_5 \in M)[\text{AREA}(x, x_1, x_2) = \text{AREA}(x_3, x_4, x_5)]$.

f maps an element of $X - M$ to **reason** $x \notin M$.

$f : X - M \rightarrow \binom{M}{2} \times \binom{M}{2} \cup \binom{M}{2} \times \binom{M}{3}$.

What is $f^{-1}(\{\{x_1, x_2\}, \{x_1, x_3\}\})$? THIS IS HARD!

What is $f^{-1}(\{x_1, x_2\}, \{x_3, x_4\})$? THIS IS HARD!

What is $f^{-1}(\{x_1, x_2\}, \{x_3, x_4, x_5\})$?

Area- $d = 3$ Case

Thm: For all $X \subseteq \mathbb{R}^3$ of size n , no three colinear, there exists Area-rainbow set of size $\Omega(n^\delta)$. (δ TBD)

Proof: Let M be a **MAXIMAL AREA-RAINBOW SET**.

Let $x \in X - M$. WHY IS x NOT IN M !? Either

- ▶ $(\exists x_1, x_2, x_3 \in M)[\text{AREA}(x, x_1, x_2) = \text{AREA}(x, x_1, x_3)]$.
- ▶ $(\exists x_1, x_2, x_3, x_4 \in M)[\text{AREA}(x, x_1, x_2) = \text{AREA}(x, x_3, x_4)]$.
- ▶ $(\exists x_1, x_2, x_3, x_4, x_5 \in M)[\text{AREA}(x, x_1, x_2) = \text{AREA}(x_3, x_4, x_5)]$.

f maps an element of $X - M$ to **reason** $x \notin M$.

$$f : X - M \rightarrow \binom{M}{2} \times \binom{M}{2} \cup \binom{M}{2} \times \binom{M}{3}.$$

What is $f^{-1}(\{\{x_1, x_2\}, \{x_1, x_3\}\})$? THIS IS HARD!

What is $f^{-1}(\{x_1, x_2\}, \{x_3, x_4\})$? THIS IS HARD!

What is $f^{-1}(\{x_1, x_2\}, \{x_3, x_4, x_5\})$? THIS IS HARD!

Area- $d = 3$ Case

Thm: For all $X \subseteq \mathbb{R}^3$ of size n , no three colinear, there exists Area-rainbow set of size $\Omega(n^\delta)$. (δ TBD)

Proof: Let M be a **MAXIMAL AREA-RAINBOW SET**.

Let $x \in X - M$. WHY IS x NOT IN M !? Either

- ▶ $(\exists x_1, x_2, x_3 \in M)[\text{AREA}(x, x_1, x_2) = \text{AREA}(x, x_1, x_3)]$.
- ▶ $(\exists x_1, x_2, x_3, x_4 \in M)[\text{AREA}(x, x_1, x_2) = \text{AREA}(x, x_3, x_4)]$.
- ▶ $(\exists x_1, x_2, x_3, x_4, x_5 \in M)[\text{AREA}(x, x_1, x_2) = \text{AREA}(x_3, x_4, x_5)]$.

f maps an element of $X - M$ to **reason** $x \notin M$.

$$f : X - M \rightarrow \binom{M}{2} \times \binom{M}{2} \cup \binom{M}{2} \times \binom{M}{3}.$$

What is $f^{-1}(\{\{x_1, x_2\}, \{x_1, x_3\}\})$? THIS IS HARD!

What is $f^{-1}(\{x_1, x_2\}, \{x_3, x_4\})$? THIS IS HARD!

What is $f^{-1}(\{x_1, x_2\}, \{x_3, x_4, x_5\})$? THIS IS HARD!

What to do?

WHAT CHANGED?

Why is this proof harder?

KEY statement about prior proof:

WHAT CHANGED?

Why is this proof harder?

KEY statement about prior proof:

1. If INVERSE IMG's are all finite so M is large.

WHAT CHANGED?

Why is this proof harder?

KEY statement about prior proof:

1. If INVERSE IMG's are all finite so M is large.
2. If INVERSE IMG's are subsets of \mathbb{R}^d or \mathbb{S}^d then induct.

WHAT CHANGED?

Why is this proof harder?

KEY statement about prior proof:

1. If INVERSE IMG's are all finite so M is large.
2. If INVERSE IMG's are subsets of \mathbb{R}^d or \mathbb{S}^d then induct.

KEY: We cared about $X \subseteq \mathbb{R}^d$ but had to work with \mathbb{S}^d as well.
NOW we will have to work with more complicated objects.

What Do Inverse Images Look Like?

$$\{x : \text{AREA}(x, x_1, x_2) = \text{AREA}(x, x_3, x_4)\} =$$

$$\{x : |\text{DET}(x, x_1, x_2)| = |\text{DET}(x, x_3, x_4)|\}.$$

What Do Inverse Images Look Like?

$$\{x : \text{AREA}(x, x_1, x_2) = \text{AREA}(x, x_3, x_4)\} =$$

$$\{x : |\text{DET}(x, x_1, x_2)| = |\text{DET}(x, x_3, x_4)|\}.$$

Def: (Informally) An **Algebraic Variety in \mathbb{R}^d** is a set of points in \mathbb{R}^d that satisfy a polynomial equation in d variables.

General Thm

Thm Let $2 \leq a \leq d + 1$. Let $r \in \mathbb{N}$. For all varieties V of dim d and degree r for all sets of n points on V there exists an a -rainbow set of size $\Omega(n^{1/(2^a-1)d})$.

General Thm

Thm Let $2 \leq a \leq d + 1$. Let $r \in \mathbb{N}$. For all varieties V of dim d and degree r for all sets of n points on V there exists an a -rainbow set of size $\Omega(n^{1/(2a-1)d})$.

Corollary Let $2 \leq a \leq d + 1$. For all $X \subseteq \mathbb{R}^d$ of size n there exists an a -rainbow set of size $\Omega(n^{1/(2a-1)d})$.

General Thm

Thm Let $2 \leq a \leq d + 1$. Let $r \in \mathbb{N}$. For all varieties V of dim d and degree r for all sets of n points on V there exists an a -rainbow set of size $\Omega(n^{1/(2a-1)d})$.

Corollary Let $2 \leq a \leq d + 1$. For all $X \subseteq \mathbb{R}^d$ of size n there exists an a -rainbow set of size $\Omega(n^{1/(2a-1)d})$.

Corollary For all $X \subseteq \mathbb{R}^d$ of size n there exists a 2-rainbow set (dist. distances) of size $\Omega(n^{1/3d})$.

General Thm

Thm Let $2 \leq a \leq d + 1$. Let $r \in \mathbb{N}$. For all varieties V of dim d and degree r for all sets of n points on V there exists an a -rainbow set of size $\Omega(n^{1/(2a-1)d})$.

Corollary Let $2 \leq a \leq d + 1$. For all $X \subseteq \mathbb{R}^d$ of size n there exists an a -rainbow set of size $\Omega(n^{1/(2a-1)d})$.

Corollary For all $X \subseteq \mathbb{R}^d$ of size n there exists a 2-rainbow set (dist. distances) of size $\Omega(n^{1/3d})$.

Corollary For all $X \subseteq \mathbb{R}^d$ of size n there is a 3-rainbow set (dist. areas) of size $\Omega(n^{1/5d})$.

General Thm

Thm Let $2 \leq a \leq d + 1$. Let $r \in \mathbb{N}$. For all varieties V of dim d and degree r for all sets of n points on V there exists an a -rainbow set of size $\Omega(n^{1/(2a-1)d})$.

Corollary Let $2 \leq a \leq d + 1$. For all $X \subseteq \mathbb{R}^d$ of size n there exists an a -rainbow set of size $\Omega(n^{1/(2a-1)d})$.

Corollary For all $X \subseteq \mathbb{R}^d$ of size n there exists a 2-rainbow set (dist. distances) of size $\Omega(n^{1/3d})$.

Corollary For all $X \subseteq \mathbb{R}^d$ of size n there is a 3-rainbow set (dist. areas) of size $\Omega(n^{1/5d})$.

Corollary For all $X \subseteq \mathbb{R}^d$ of size n there is a 4-rainbow set (dist. volumes) of size $\Omega(n^{1/7d})$.

Comments on the Proof

General Thm

Thm Let $2 \leq a \leq d + 1$. Let $r \in \mathbb{N}$. For all varieties V of dim d and degree r for all sets of n points on V there exists an a -rainbow set of size $\Omega(n^{1/(2a-1)d})$.

Corollary Let $2 \leq a \leq d + 1$. For all $X \subseteq \mathbb{R}^d$ of size n there exists an a -rainbow set of size $\Omega(n^{1/(2a-1)d})$.

Corollary For all $X \subseteq \mathbb{R}^d$ of size n there exists a 2-rainbow set (dist. distances) of size $\Omega(n^{1/3d})$.

Corollary For all $X \subseteq \mathbb{R}^d$ of size n there is a 3-rainbow set (dist. areas) of size $\Omega(n^{1/5d})$.

Corollary For all $X \subseteq \mathbb{R}^d$ of size n there is a 4-rainbow set (dist. volumes) of size $\Omega(n^{1/7d})$.

Comments on the Proof

1. Proof uses Algebraic Geometry in Proj Space over \mathbb{C} .

General Thm

Thm Let $2 \leq a \leq d + 1$. Let $r \in \mathbb{N}$. For all varieties V of dim d and degree r for all sets of n points on V there exists an a -rainbow set of size $\Omega(n^{1/(2a-1)d})$.

Corollary Let $2 \leq a \leq d + 1$. For all $X \subseteq \mathbb{R}^d$ of size n there exists an a -rainbow set of size $\Omega(n^{1/(2a-1)d})$.

Corollary For all $X \subseteq \mathbb{R}^d$ of size n there exists a 2-rainbow set (dist. distances) of size $\Omega(n^{1/3d})$.

Corollary For all $X \subseteq \mathbb{R}^d$ of size n there is a 3-rainbow set (dist. areas) of size $\Omega(n^{1/5d})$.

Corollary For all $X \subseteq \mathbb{R}^d$ of size n there is a 4-rainbow set (dist. volumes) of size $\Omega(n^{1/7d})$.

Comments on the Proof

1. Proof uses Algebraic Geometry in Proj Space over \mathbb{C} .
2. Proof uses Maximal subsets in same way as easier proofs.

General Thm

Thm Let $2 \leq a \leq d + 1$. Let $r \in \mathbb{N}$. For all varieties V of dim d and degree r for all sets of n points on V there exists an a -rainbow set of size $\Omega(n^{1/(2a-1)d})$.

Corollary Let $2 \leq a \leq d + 1$. For all $X \subseteq \mathbb{R}^d$ of size n there exists an a -rainbow set of size $\Omega(n^{1/(2a-1)d})$.

Corollary For all $X \subseteq \mathbb{R}^d$ of size n there exists a 2-rainbow set (dist. distances) of size $\Omega(n^{1/3d})$.

Corollary For all $X \subseteq \mathbb{R}^d$ of size n there is a 3-rainbow set (dist. areas) of size $\Omega(n^{1/5d})$.

Corollary For all $X \subseteq \mathbb{R}^d$ of size n there is a 4-rainbow set (dist. volumes) of size $\Omega(n^{1/7d})$.

Comments on the Proof

1. Proof uses Algebraic Geometry in Proj Space over \mathbb{C} .
2. Proof uses Maximal subsets in same way as easier proofs.
3. Proof is by induction on d .

Open Questions

Open Questions

1. Better Particular Results: e.g., want
for all $X \subseteq \mathbb{R}^2$ of size n , there exists a rainbow set of size $\Omega(n^{1/2})$.

Open Questions

1. Better Particular Results: e.g., want
for all $X \subseteq \mathbb{R}^2$ of size n , there exists a rainbow set of size $\Omega(n^{1/2})$.
2. General Better Results: e.g., want
Let $1 \leq a \leq d + 1$. For all $X \subseteq \mathbb{R}^d$ of size n there exists a rainbow set of size $\Omega(n^{1/ad})$.

Open Questions

1. Better Particular Results: e.g., want
for all $X \subseteq \mathbb{R}^2$ of size n , there exists a rainbow set of size $\Omega(n^{1/2})$.
2. General Better Results: e.g., want
Let $1 \leq a \leq d + 1$. For all $X \subseteq \mathbb{R}^d$ of size n there exists a rainbow set of size $\Omega(n^{1/ad})$.
3. Get easier proofs of general theorem.

Open Questions

1. Better Particular Results: e.g., want
for all $X \subseteq \mathbb{R}^2$ of size n , there exists a rainbow set of size $\Omega(n^{1/2})$.
2. General Better Results: e.g., want
Let $1 \leq a \leq d + 1$. For all $X \subseteq \mathbb{R}^d$ of size n there exists a rainbow set of size $\Omega(n^{1/ad})$.
3. Get easier proofs of general theorem.
4. Find any nontrivial limits on what we can do. (Trivial: $n^{1/d}$).

Open Questions

1. Better Particular Results: e.g., want
for all $X \subseteq \mathbb{R}^2$ of size n , there exists a rainbow set of size $\Omega(n^{1/2})$.
2. General Better Results: e.g., want
Let $1 \leq a \leq d + 1$. For all $X \subseteq \mathbb{R}^d$ of size n there exists a rainbow set of size $\Omega(n^{1/ad})$.
3. Get easier proofs of general theorem.
4. Find any nontrivial limits on what we can do. (Trivial: $n^{1/d}$).
5. Algorithmic aspects.